

PSMSL Report to IAG – April 2013

*Lesley Rickards, Director, Permanent Service for Mean Sea Level (PSMSL)
National Oceanography Centre, Joseph Proudman Building, 6 Brownlow Street, Liverpool L3 5DA, UK*

1. Introduction

Since 1933, the Permanent Service for Mean Sea Level (PSMSL) has operated at the National Oceanography Centre (NOC), Liverpool (previously the Proudman Oceanographic Laboratory), with the aims of providing the global data bank for long term sea level information from tide gauges, and of providing a wider Service to the sea level community. It was a member of the Federation of Astronomical and Geophysical Data Analysis Services (FAGS) until its dissolution and works under the auspices of the International Council for Science (ICSU) and is applying for membership of the new ICSU World Data System (WDS).

The data set and ancillary information are provided free of charge and are made available to the international scientific community through the PSMSL website. The metadata includes descriptions of benchmarks and their locations, types of instrumentation and frequency of data collection (where available) as well as notes on other issues that we feel the users should be aware of (e.g. earthquakes that are known to have occurred in the vicinity or subsidence due to local groundwater extraction). The free access to data by users is central to the PSMSL's mission, and conversely no supplier is ever paid for their data, nor are licensing terms ever entered into.

2. PSMSL-related scientific meetings, activities and events

PSMSL staff have continued to be active participants in the IOC Group of Experts on the Global Sea Level Observing System (GLOSS) and GGOS meetings, co-convened sea level sessions at the EGU and contributed to IOC coordination group tsunami warning system meetings. PSMSL has contributed to the IPCC Fifth Assessment Report with Dr Svetlana Jevrejeva a lead author for Working Group I and other PSMSL staff also contributing.

2013 marks the 80th anniversary of the foundation of the PSMSL. To commemorate this PSMSL is hosting or co-convening the following events:

- A workshop in Liverpool, UK, on major research topics in sea level science. The workshop, to be held in October, will include talks reviewing aspects of the IPCC Fifth Assessment Report (Working Group I). There will also be presentations covering many aspects of regional variability in sea level.
- A symposium entitled "Implications of sea level change for the coastal zone - A symposium to mark the 80th anniversary of the Permanent Service for Mean Sea Level (PSMSL)" at the IAHS/IASO/IASPEI Joint Assembly in Gothenburg, Sweden (July 2013).
- A session at EGU 2013: Global and regional sea level rise and variability: from past to future.

3. Collection, analysis, publication and interpretation of monthly and annual means of sea level from the global network of tide gauges

Since the start of 2012, approximately 16468 station-months of MSL data from about 700 stations were added to the PSMSL databank (and a further 976 months were updated), bringing the total PSMSL data holdings to over 707402 station-months from 2174 stations. Most of the data originated from Europe and North America together with significant data sets from Japan and Australia. There are gaps in data receipts from parts of SE Asia, central and South America; these are presently being targeted to try to improve data flow. Africa continues to receive special attention through ODINAfrica and IOTWS, although data flow has improved considerably over the last decade. Close links have been maintained with the University of Hawaii Sea Level Center and other international sea level data centres.




Figure 1a: Number of station years added to PSMSL database in 2012

Figure 1b: Number of station with data added to PSMSL database in 2012




Figure 2: Distribution of all PSMSL stations, both research quality (RLR) and non-datum controlled stations (known as ‘Metric only’)

3.1 Interactive map showing long-term trends

The relative sea level trends map allows interactive investigation of global mean sea level trends since 1900. The limits of the period to be viewed can be selected by either moving the buttons on the slider or by entering the values in the two text boxes. A period of at least thirty years must be selected. The map will display the annual sea level trend at each station that has suitable data available over the selected period. Further information is available on the methods page (see: <http://www.psmsl.org/products/trends/methods.php>).

Note that these measured trends are not corrected for local land movement. Furthermore, no attempt has been made to assess the validity of any individual fit, so results should not be treated as a publication quality values suitable for use in planning or policy making.

The map should be used with some care as anomalous trends have many causes:

- land movements (e.g. earthquakes, glacial isostatic adjustment)
- unexplained instrumental datum shifts
- changes in atmospheric pressure
- short records

A more complete account can be found in the geophysical signals section of the PSMSL web-site (see: http://www.psmsl.org/train_and_info/geo_signals/). A table of long term trends derived from annual mean values of sea level in the PSMSL RLR data set demonstrates the rate of change of sea level at each station.





Figure 3: Sample map showing relative sea level trends

3.2 Interactive map showing sea level anomalies

Annual mean sea level can vary considerably from year to year in response to various meteorological and oceanographic forcings, typically by hundreds of millimetres. The product allows one to examine the global variations in a year of your choice: select this year using either the slider or the text box. The map presents the difference between the annual RLR data for each station (which is quality and datum controlled) compared to that station's long term mean over the baseline period of 1960-1990.

The long term trend at each station (estimated using the baseline period) can be removed if required. This will prevent results being dominated by long term changes, but will result in the loss of stations for which there is not enough data to calculate a trend. Further information is provided on the methods and derived trends pages of the PSMSL web-site.



**Figure 4: Sea level anomalies for 2010 relative to 1960-1990.
(Top image: not detrended. Bottom image: detrended)**

4. Summary and forward look

It can be seen that PSMSL continues to be active with regard to workshops/conferences and with data acquisition and analysis. The functions provided by the PSMSL are in as much demand as ever and new products continue to be developed. Future plans include:

- Improved integration of the mean sea level data set with higher frequency data and improving the quality of accompanying metadata;
- Keeping contact with data suppliers (the trend being to acquire data from websites rather than direct supply) and ensuring that data made available in real-time are also contributed to PSMSL;
- Inclusion of information on uncertainties/errors in the tide gauge data;
- Addition of bottom pressure record section and data to the PSMSL web-site;
- Redevelopment of capacity building/training material.

Particular thanks as usual go to PSMSL staff and to colleagues at the National Oceanography Centre and British Oceanographic Data Centre who contribute part of their time to PSMSL activities.